

E-HEALTH-COM, Competence, Heft 2/2009

Präventive Bestimmung des Herz-Kreislauf-Risikos anhand der photoplethysmographisch gemessenen Pulswelle

Autoren

Ludger Grote

Sleep Laboratory
Department of Pulmonary Medicine
Sahlgrenska University Hospital
SE-41315 Göteborg

Dirk Sommermeyer

MCC Gesellschaft für Diagnosesysteme in Medizin und Technik mbH & Co.KG
76135 Karlsruhe
dirk.sommermeyer@mcc-med.de

Literatur

- [1] Statistisches Bundesamt, Wiesbaden, 2005
- [2] Paul E. Peppard, Ph.D., Terry Young, Ph.D., Mari Palta, Ph.D., James Skatrud, M.D. Prospective Study of the Association between Sleep-Disordered Breathing and Hypertension NEJM, 2000; 342: 1378-1384
- [3] European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. J Hypertens 2003; 21(6):1011-1053
- [4] Mallat S, Zhang Z. Matching Pursuit in a timefrequency dictionary. IEEE Transactions on Signal Processing 1993; 41:3397-3415
- [5] Wang XH, Hong JR. On the handling of fuzziness for continuous-valued attributes in decision tree generation. Fuzzy Sets and Systems 1998; 99(283-290)